


MOLECULAR CELL BIOLOGY

JAMES DARNELL

*Vincent Astor Professor
Rockefeller University*


HARVEY LODISH

*Member of the Whitehead Institute for
Biomedical Research
Professor of Biology, Massachusetts
Institute of Technology*



DAVID BALTIMORE

*Director of the Whitehead Institute for
Biomedical Research
Professor of Biology, Massachusetts
Institute of Technology*



BEST AVAILABLE COPY

**SCIENTIFIC
AMERICAN
BOOKS**

Applicants: Bernard F. Erlanger et al.
U.S. Serial No.: 10/618,179
Filed: July 11, 2003
Exhibit A

Front cover photograph courtesy of J. Victor Small

Back cover photograph courtesy of Richard Feldman

Library of Congress Cataloging-in-Publication Data

Darnell, James E.
Molecular cell biology.

Includes bibliographies and index.

1. Cytology. 2. Molecular biology. I. Lodish, Harvey F. II. Baltimore, David. III. Title.
[DNLM: 1. Cells. 2. Molecular Biology. QH 581.2 D223m]
QH581.2.D37 1986 574.87'6042 86-1881
ISBN 0-7167-1448-5
ISBN 0-7167-6001-0 (international student ed.)

Copyright © 1986 by Scientific American Books, Inc.

No part of this book may be reproduced by any mechanical, photographic, or electronic process, or in the form of a phonographic recording, nor may it be stored in a retrieval system, transmitted, or otherwise copied for public or private use, without the written permission of the publisher.

Printed in the United States of America

Scientific American Books is a subsidiary of Scientific American, Inc.
Distributed by W. H. Freeman and Company, 41 Madison Avenue,
New York, New York 10010


BEST AVAILABLE COPY

making a total of eight types of immunoglobulin. Attached to all the H chains are asparagine-linked carbohydrate chains.

Every individual antibody molecule has one type of L chain and one type of H chain. The chains are held together by disulfide bonds to form a monomer, and two monomers are linked by disulfide bonds to form the basic dimeric structure of the molecule (Figure 24-15). Within each chain, units made of about 110 amino acids fold up to form compact *domains*. Each domain has a single internal disulfide bond which holds it together. An L chain has two domains, H chains have four or five domains. The first two N-terminal domains of the H chains interact with the two L-chain domains, producing a compact unit that acts as the binding region of the antibody. In most H chains, a *hinge* region consisting of a small number of

amino acids is found after the first two domains. The hinge is flexible and allows the binding regions to move freely relative to the rest of the molecule. At the hinge region are located the cysteine residues whose SH groups are linked to form the —S—S— bridges between the two monomer units of the antibody dimer. The hinge region are the places most susceptible on the molecule to the action of protease; light protease treatment can split an antibody into two pieces, called F_{ab} and F_c fragments. The F_{ab} portion has the antigen-binding site, the F_c portion has the effector regions (see Figure 24-15).

The very first amino acid sequences determined on H chains from human myelomas made it clear that the N-terminal domain has a very variable structure while the C-terminal domain has a quite constant structure. The N-terminal domain is called the *variable region* and the C-terminal domain is called the *constant region* (Figure 24-15). H-chain sequences show the same diversity within each class the N-terminal domain is highly variable and the C-terminal domains have a constant sequence. The variable domains of L and H chains are bound to one another. In fact, they interact closely to form a single compact unit (Figure 24-15). This unit is the antibody-binding site, the region of the antibody molecule that binds to antigen. This can be demonstrated by the use of an antigen that contains a reactive chemical group: the reactive group on the antigen will form a covalent bond to the variable domain of the H or L chain, showing that the variable domains form the antigen-binding pocket.



Domain structure and the complementarity-determining regions (CDRs) of an antibody. (a) The molecule is organized in disulfide-bonded 110 amino acid domains, four in the heavy chain and two in the light chain. The farthest N-terminal domain of each chain is variable in sequence (V region); the other domains are constant in sequence (regions C_{H1} , C_{H2} , and C_{H3} in the heavy chain and C_L in the light chain). When the molecule is protease-digested, cutting of the hinge region connecting C_{H1} and C_{H2} , the most sensitive spot on the molecule, splits the molecule into two parts, F_{ab} (the antigen-binding domain) and F_c (the effector region). Within the V region are segments of highly variable sequence containing the CDRs, the amino acids that actually contact the antigen. (b) Antibody binding site. By the coordinate effort of the CDRs from light and heavy chains, a binding pocket is formed into which a specific antigen fits.

BEST AVAILABLE COPY